



The
Patent
Office

REC'D

OCT 1998

WIPO

PCT

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP9 1RH

PCT/GB 98 / 03056

S

09/509863

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed

A. Brewster

Dated 2 October 1998

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Request for grant of a patent

(See the notes on the back of this form. You can also get a explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road
Newport
Gwent NP9 1RH

1. Your reference C97/082F

2. Patent application number
(The Patent Office will fill in this part) **9721588.3**

10 OCT 1997

3. Full name, address and postcode of the or of each applicant (underline all surnames)
E. I. DuPont de Nemours and Company
1007 Market Street
Wilimington
Delaware 198 98
UNITED STATES OF AMERICA

Patents ADP number (if you know it) 730 1484001

If the applicant is a corporate body, give the country/state of its incorporation USA

4. Title of the invention TEXTILE TREATMENT

5. Name of your agent (if you have one) MEWBURN ELLIS

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)
YORK HOUSE
23 KINGSWAY
LONDON
WC2B 6HP

Patents ADP number (if you know it) 109006

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number	Country	Priority application number (if you know it)	Date of filing (day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application	Number of earlier application	Date of filing (day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:
a) any applicant named in part 3 is not an inventor, or
b) there is an inventor who is not named as an applicant, or
c) any named applicant is a corporate body.
See note (d)) YES

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form 0

Description 30

Claim(s) 4

Abstract 0

Drawing(s) 0

10. If you are also filing any of the following, state

Priority documents 0

Translations of priority documents 0

Statement of inventorship and right to grant of a patent (Patents Form 7/77) 0

Request for preliminary examination and search (Patents Form 9/77) 0

Request for substantive examination (Patents Form 10/77) 0

Any other documents 0
(Please specify)

11. I/We request the grant of a patent on the basis of this application.

Signature
Michael Ford

Date
10 October 1997

12. Name and daytime telephone number of person to contact in the United Kingdom MICHAEL F. FORD 0171 240 4405

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- a) If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- b) Write your answers in capital letters using black ink or you may type them.
- c) If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- d) If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- e) Once you have filled in the form you must remember to sign and date it.
- f) For details of the fee and ways to pay please contact the Patent Office.

Textile Treatment

This invention relates to the treatment of textiles containing spandex fibres.

At the present time, many garments are made from fabric which contains a mixture of fibres, a proportion of which are elastic, so that the fabric has the ability to stretch and to recover from stretch. Spandex fibres are commonly used for this purpose. The term "Spandex" has been adopted as a generic term by the United States Federal Trade Commission to denote a manufactured fibre in which the fibre-forming substance is a long chain synthetic polymer composed of at least 85% of a segmented polyurethane. A discussion of such fibres can be found in "History of Spandex Elastomeric Fibres" by A.J. Ultee, which is a chapter starting at page 278 in *Man-Made Fibres: Their Origin and Development*, edited by R.V. Seymour and R.S. Porter, Elsevier 1993. Spandex fibres are also referred to as "elastane" or "elasthane" fibres.

Another discussion of such fibres is found under the heading "Segmented Polyurethanes" at page 613 of *Handbook of Textile Fibres* by J. Gordon Cook, 5th Ed. Merrow Publishing Company 1984. Further description of elastanes and their applications can be found in "Synthesefasern: Grundlagen, Technologie, Verarbeitung und Anwendung", B von Falkei (editor), Verlag Chemie (1981). Commercially

available elastanes are well known, in particular as sold under the name LYCRA®, a registered trade mark of DuPont de Nemours and Company. Patents relating to such fibres include US-A-5000899, US-A-5288779 and US-A-5362432.

5 The deposition of perfume onto garments and other fabrics during laundering has been established for many years. Perfume is incorporated into laundry products such as detergent compositions for fabric washing and rinse conditioners for softening the fabrics.

10 Although the perfume serves to cover the base odour of such a product and to give the unused product an attractive fragrance, it also deposits on the fabric.

Certain perfumes have the ability to provide deodorant action against body odour, either when directly applied to human skin, or when included in a laundry product. Such
15 perfumes are described in EP-B-3172, US-A-4304679, US-A-4278658, US-A-4134838, US-A-4288341 and US-A-4289641, US-A-5482635 and US-A-5554588.

Summary of the Invention

20 We have now found that a number of fragrance materials used in perfumery are able to deposit and then be retained better on spandex fibres than on a number of other textile fibres.

Delivery of fragrance materials to fabric can take place during washing, as is well known. The present invention appreciates that the application of perfume to textiles containing spandex fibres can be utilised in the treatment of textiles which are newly made - that is to say textile goods which have never been worn as garments by a consumer.

Therefore in one aspect the present invention provides a method of treating textile which is yarn or unworn fabric containing spandex fibres, comprising contacting the textile with a perfume so that the perfume is deposited on the fabric.

In a related second aspect, the invention provides textile which is yarn or unworn fabric containing spandex fibres, having perfume deposited on the textile.

We have observed that a range of fragrance materials deposited on such textiles will still be perceptible on the fabric, even after several washes of the fabric using laundry products with a different perfume, or none.

The invention also provides use of a perfume composition in the treatment of textile which is yarn or unworn fabric containing spandex and other fibres, to deposit fragrance materials at a greater concentration on the spandex fibres than on the other fibres.

In significant forms of this invention, the perfume used to treat the textile (or the combination of fragrance materials deposited thereon) is a deodorant perfume. Then when the textile is made into a garment, that garment will have an in-built deodorant property.

5 Detailed description

The various aspects of this invention, preferred forms and materials useful therein will now be discussed in greater detail.

Textiles

- 10 The textiles to which this invention relates include spandex fibres. As mentioned earlier, this term denotes a manufactured fibre in which the fibre forming substance is a long chain synthetic polymer compound composed of at least 85% of a segmented polyurethane.
- 15 Thus the polymer which is spun into spandex fibres is a copolymer incorporating urethane linkages. Generally the polymer contains so-called soft (i.e. lower melting) segments which may be polyalkylene ethers or polyesters and
- 20 so-called hard (i.e. higher melting) segments which are portions derived from the reaction of an isocyanate and a chain extender which is typically a diamine.

The soft segments may be poly(tetramethylene)ethers, possibly containing substituted tetramethylene glycol residues as described in US-A-5000899. Organic diisocyanates which may be used include conventional diisocyanates, such as diphenylmethane-4,4'-diisocyanate, also known as methylene-bis(4-phenylisocyanate) or "MDI", 2,4-tolylene diisocyanate, methylene-bis(4-cyclohexylisocyanate), isophorone diisocyanate, tetramethylene-p-xylylene diisocyanate, and the like. MDI is preferred.

- 10 Chain extenders used in producing the hard segment of the fibres preferably include one or more of ethylenediamine (EDA), 1,3-propylenediamine, 1,4-cyclohexanediamine, hydrogenated m-phenylenediamine (HPMD), 2-methylpentamethylene diamine (MPMD) and 1,2-propylene
- 15 diamine. More preferably, the chain extender is one or more of ethylenediamine, 1,3-propylenediamine, and 1,4-cyclohexanediamine, optionally mixed with HPMD, MPMD and/or 1,2-propylenediamine.

- Spandex fibres with poly(tetramethylene)ethers as the soft
- 20 segments are marketed by DuPont de Nemours International S.A. under the registered trade mark LYCRA[®] of DuPont de Nemours and Company.

Spandex fibres are generally mixed with other fibres such as cotton, polyamide, wool, polyester and acrylics and made

into yarn which is then made into fabric. The contents of spandex fibres is usually in a range from 0.5% by weight of the yarn or fabric up to 50%, more usually from 1 to 30% by weight of the yarn of fabric.

5 A wide range of garments may contain spandex fibres in the fabric, including active sports wear, intimate apparel, hosiery and a variety of ready to wear casual clothing.

The textiles which are treated with a perfume composition prior to wearing may be yarn which is later made into fabric, or may be fabric in the form of a web or lengths
10 from a web which have not yet been made into garments, or may possibly be garments which have been made but not yet been worn.

Preferably the treatment with a perfume composition is carried out while treating with other material in a
15 conventional process step, especially a wet stage in which the yarn or fabric is treated with a finishing agent to improve its hand or appearance.

The materials which may be applied to fabric in a conventional finishing treatment include resins to confer
20 stiffness, fabric stability or permanent press, fabric softeners, flame retardants, fabric brighteners, anti-sag agents, materials to confer soil or stain resistance and water repellants.

Techniques which are conventionally used to apply such materials are padding and exhaustion, both well known in the technology of textile manufacture.

Treatment with a perfume composition in accordance with this invention can be carried out by including the perfume
5 composition in the liquor used in a process as above.

The amount of perfume deposited on the fabric in a treatment step carried out on unworn fabric will generally be from 0.001% to 1% by weight of the fabric.

Fragrance materials

10 We have found that a range of fragrance materials deposit well on, or are retained well on, spandex fibres. Such materials include the following two categories:

Category A

15 hydroxylic materials which are alcohols, phenols or salicylates, with an octanol/water partition coefficient (P) whose common logarithm ($\log_{10}P$) is 2.5 or greater, and a gas chromatographic Kovats index (as determined on polydimethylsiloxane as non-polar stationary phase) of at least 1050.

Category B

esters, ethers, nitriles, ketones or aldehydes; with an octanol/water partition coefficient (P) whose common logarithm ($\log_{10}P$) is 2.5 or greater, and a gas chromatographic Kovats index (as determined on
5 polydimethylsiloxane as non-polar stationary phase) of at least 1300.

The octanol-water partition coefficient (or its common logarithm 'logP') is well known in the literature as an indicator of hydrophobicity and water solubility (see
10 Hansch and Leo, *Chemical Reviews*, 71, 526-616, (1971); Hansch, Quinlan and Lawrence, *J.Organic Chemistry*, 33, 347-350 (1968). Where such values are not available in the literature they may be measured directly, or estimated approximately using mathematical algorithms. Software
15 providing such estimations are available commercially, for example 'LogP' from Advanced Chemistry Design Inc.

A requirement for $\log_{10}P$ of 2.5 or more calls for materials which are somewhat hydrophobic.

Kovats indices are calculated from the retention time in a
20 gas chromatographic measurement referenced to the retention time for alkanes [see Kovats, *Helv.Chim.Acta* 41, 1915 (1958)]. Indices based on the use of a non-polar stationary phase have been used in the perfumery industry for some

years as a descriptor relating to the molecular size and boiling point of ingredients. A review of Kovats indices in the perfume industry is given by T Shibamoto in "Capillary Gas Chromatography in Essential Oil Analysis", P Sandra and C Bicchi (editors), *Huethig* (1987), pages 259-274. A common
5 non-polar phase which is suitable is 100% dimethyl polysiloxane, as supplied for example under a variety of tradenames such as HP-1 (Hewlett-Packard), CP Sil 5 CB (Chrompack), OV-1 (Ohio Valley) and Rtx-1 (Restek).

Materials of low Kovats index tend to be volatile and are
10 not retained well on many fibres.

We have found that when perfumery materials have partition coefficient as above and a relatively high value of Kovats index, deposition and retention on spandex tends to be greater than on other fibres. Preferably therefore, the
15 perfume composition contains at least 50 wt %, better at least 70 or 80 wt % of materials from the categories above.

We have found that there is a particularly high enhancement of deposition and retention on spandex, compared to other fibres, with materials within the above categories and
20 having a Kovats index of not more than 1600. These sub-sets of categories A and B may be termed categories A' and B'. Preferably therefore, the perfume composition contains at least 10 wt%, better at least 20 wt% or 25 wt% of such materials. In some preferred perfumes the amount of

fragrance materials from categories A' and B' is at least 40 wt% in total.

Such fragrance materials are of mid-range volatility (i.e. intermediate between the volatile perfume materials used as "top-notes" and the materials of low volatility which are customarily used as base notes in perfumes). These materials of mid-range volatility are often not perceptible on other fabrics such as cotton, polyamide and polyester after washing and drying.

Category A includes alcohols of general formula ROH where the hydroxyl group may be primary, secondary or tertiary, and the R group is an alkyl or alkenyl group, optionally branched or substituted, cyclic or acyclic, such that ROH has partition coefficient and Kovats properties as defined above. Alcohols of Kovats index 1050 to 1600 are typically monofunctional alkyl or arylalkyl alcohols with molecular weight falling within the range 150 to 230.

Category A also includes phenols of general formula ArOH, where the Ar group denotes a benzene ring which may be substituted with one or more alkyl or alkenyl groups, or with an ester grouping $-CO_2A$, where A is a hydrocarbon radical, in which case the compound is a salicylate. ArOH has partition coefficient and Kovats index as defined above. Typically, such phenols with Kovats index 1050 to 1600 are monohydroxylic phenols with molecular weight

falling within the range 150 to 210.

A sub-set of fragrance materials which are particularly preferred are those with a partition coefficient of 1000 or more, i.e. $\log_{10}P$ of 3 or more, and a Kovats parameter of 1100 up to 1600.

- 5 Some examples of hydroxylic ingredients which fulfil the above criteria for category A' are listed as a table below. Materials which are in the preferred sub-set are marked with an asterisk. Semitrivial names are those used in standard texts known within the perfume industry,
- 10 particularly: *Common Fragrance and Flavor Materials* by Bauer, Garbe and Surburg, VCH Publ., 2nd edition (1990), and *Perfume and Flavour Materials*, Steffen Arctander, published in two volumes by the author (1969).

Examples of fragrance materials in category A'	
15	1-(2'-tert-butylcyclohexyloxy)-butan-2-ol*
	3-methyl-5-(2',2',3'-trimethylcyclopent-3-enyl)-pentan-2-ol*
	4-methyl-3-decen-5-ol*
	amyl salicylate*
20	2-ethyl-4(2',2',3-trimethylcyclopent-3'-enyl)but-2-enol* (Banalol, TM)
	borneol*
	carvacrol*
	citronellol*
25	9-decenol*
	dihydroeugenol*
	dihydrolinalol*

	dihydromyrcenol	
	dihydroterpineol *	
	eugenol	
	geraniol*	
	hydroxycitronellal*	
5	isoamyl salicylate*	
	isobutyl salicylate*	
	isoeugenol*	
	linalol	
	menthol*	
10	nerolidol*	
	nerol*	
	para tert-butyl cyclohexanol*	
	phenoxanol*	
	terpineol	
15	tetrahydrogeraniol*	
	tetrahydrolinalol	
	tetrahydromyrcenol	
	thymol*	
	2-methoxy-4-methylphenol	(Ultravanil, TM)
20	(4-isopropylcyclohexyl) -methanol*	

Some examples of fragrance materials which are in category A but which have Kovats index above 1600 (so as to fall outside category A') are:

benzyl salicylate

25 cyclohexyl salicylate

hexyl salicylate

patchouli alcohol

farnesol

Category B is esters, ketones, aldehydes, nitriles or ethers which have an octanol-water partition coefficient whose common logarithm ($\log_{10}P$) is at least 2.5, and a Kovats index of at least 1300 (non-polar phase).

Ingredients of Category B are of general formula RX, where
5 X may be in a primary, secondary or tertiary position, and is one of the following groups: -COA, -OA, -CO₂A, -CN or -CHO. The groups R and A are hydrocarbon residues, cyclic or non-cyclic and optionally substituted. In some forms of this invention, category B excludes any material with a
10 free hydroxy group, so that where a hydroxyl group is present, the material should be considered only for Category A membership. Typically, the materials of Category B with Kovats index not exceeding 1600 (which may be called category B') are monofunctional compounds with molecular
15 weights in the range 160 to 230.

A sub-set of particularly preferred fragrance materials within category B' is those with a Kovats parameter falling within the range 1350 up to 1600, and possessing a molecular structure containing a ring, such as phenyl or
20 cycloalkyl.

A number of fragrance materials which fulfil the above criteria for category B' are listed in the table below. Materials which are in the preferred sub-set are marked

with an asterisk.

Examples of fragrance materials in category B'	
	1-methyl-4-(4-methyl-3-pentenyl)-3-cyclohexene-1-carbaldehyde*
	1-(5',5'-dimethylcyclohexenyl)-pent-en-1-one*
5	2-heptyl cyclopentanone*
	2-methyl-3-(4'-tert-butylphenyl)propanal
	2-methylundecanal
	2-undecenal
	2,2-dimethyl-3-(4'-ethylphenyl)-propanal
10	3-(4'-isopropylphenyl)-2-methylpropanal
	4-methyl-4-phenylpent-2-yl acetate*
	allyl cyclohexyl propionate*
	allyl cyclohexyloxyacetate*
	amyl benzoate*
15	methyl ethyl ketone trimers (Azarbre, TM)
	benzophenone*
	3-(4'-tert-butylphenyl)-propanal (Bourgeonal, TM)
	caryophyllene*
	cis-jasmone*
20	citral diethyl acetal
	citronellal diethyl acetal
	citronellyl acetate
	phenylethyl butyl ether (Cressanther, TM)
	damascone, alpha-*
25	damascone, beta-*
	damascone, delta-*
	decalactone, gamma-*
	dihydro isojasmonate*
	dihydrojasmone*
30	dihydroterpinyl acetate
	dimethyl anthranilate*
	diphenyl oxide*

	diphenylmethane*
	dodecanal
	dodecen-2-al
	dodecane nitrile
	1-ethoxy-1-phenoxyethane (Efetaal, TM)
5	3-(1'-ethoxyethoxy)-3,7-dimethylocta-1,6-diene (Elintaal Forte (TM))
	4-(4'-methylpent-3'-enyl)-cyclohex-3-enal (Empetaal, TM)
	ethyl tricyclo[5.2.1.0~2,6~]decane-2-carboxylate*
	1-(7-isopropyl-5-methylbicyclo[2.2.2]oct-5-en-2-yl)-1-ethanone* (Felvinone, TM)
10	allyl tricyclodecenyl ether* (Fleuroxene, TM)
	tricyclodecenyl propanoate* (Florocyclene, TM)
	gamma-undecalactone*
	n-methyl-n-phenyl-2-methylbutanamide* (Gardamide, TM)
	tricyclodecenyl isobutyrate* (Gardocyclene, TM)
15	geranyl acetate
	hexyl benzoate*
	ionone alpha*
	ionone beta*
	isobutyl cinnamate*
20	isobutyl quinoline*
	isoeugenyl acetate*
	2,2,7,7-tetramethyltricycloundecan-5-one* (Isolongifolanone, TM)
	tricyclodecenyl acetate* (Jasmacyclene, TM)
25	2-hexylcyclopentanone (Jasmatone, TM)
	4-acetoxy-3-pentyltetrahydropyran* (Jasmopyrane, TM)
	ethyl 2-hexylacetoacetate (Jessate, TM)
	8-isopropyl-6-methylbicyclo[2.2.2]oct-5-ene-2-carbaldehyde (Maceal, TM)
30	methyl 4-isopropyl-1-methylbicyclo[2.2.2]oct-5-ene-2-carboxylate*
	methyl cinnamate
	alpha iso methyl ionone*
	methyl naphthyl ketone*

	nerolin	
	nonalactone gamma	
	nopyl acetate*	
	para tert-butyl cyclohexyl acetate	
5	4-isopropyl-1-methyl-2-[1'-propenyl]-benzene*	(Pelargene, TM)
	phenoxyethyl isobutyrate*	
	phenylethyl isoamyl ether*	
	phenylethyl isobutyrate*	
	tricyclodecenyl pivalate*	(Pivacyclene, TM)
10	phenylethyl pivalate*	(Pivarose, TM)
	phenylacetaldehyde hexylene glycol acetal*	
	2,4-dimethyl-4-phenyltetrahydrofuran	(Rhubafuran, TM)
	rose acetone*	
	terpinyl acetate	
15	4-isopropyl-1-methyl-2-[1'-propenyl]-benzene	(Verdoracine, TM)
	yara*	
	(4-isopropylcyclohexadienyl)ethyl formate	

Examples of fragrance materials which lie within category
 20 B, but have Kovats index above 1600 and so are outside
 category B' are listed in the following table:

	Within category B but outside category B	
	amyl cinnamate	
	amyl cinnamic aldehyde	
25	amyl cinnamic aldehyde dimethyl acetal	
	cinnamyl cinnamate	
	1,2,3,5,6,7,8,8a,-octathyro-1,2,8,8-tetramethyl-2-acetyl naphthalene	(iso E super (TM))
30	cyclo-1,13-ethylenedioxytridecan-1,13-dione	(ethylene brassylate)
	cyclopentadecanolide	(Exaltolide, TM)
	hexyl cinnamic aldehyde	

	1,3,4,6,7,8-hexahydro-4,6,6,7,8,8-hexamethylcyclopenta[g]- 2-benzopyran	(Galaxolide, TM)
	geranyl phenyl acetate	
	6-acetyl-1-isopropyl-2,3,3,5-tetramethylindane	(Traseolide, TM)
5	1,1,2,4,4,7-hexamethyl-6-acetyl-1,2,3,4- tetrahydronaphthalene	(Tonalid, TM)

As indicated above, it is particularly preferred to utilise a perfume composition which has deodorant properties.

Preferably, the perfume is a deodorant perfume giving a
10 Malodour Reduction Value of at least 0.5, preferably at least 0.9, in the Malodour Reduction Value test described below and which is an adaptation of the test described in EP-A-147191 and corresponding US-A-4663068.

The Malodour Reduction Value Test

15 In this test, the Malodour Reduction Value of a deodorant perfume is measured by assessing its effectiveness, when applied to fabric, in reducing body malodour when the fabric so treated is placed in contact with the axillae (armpits) of a panel of human subjects, and held there for
20 a standard period of time. From subsequent olfactory evaluation by trained assessors, a Malodour Reduction Value can be calculated so giving a measure of the effectiveness as a deodorant of the perfume under test.

Stage 1 is preparation of the perfume treated fabric. A
25 test fabric is subjected to a textile finishing which

applies perfume to the fabric at a predetermined percentage of perfume composition, by weight of the cloth. A control fabric is given similar treatment, with or without perfume, depending on the purpose of the test. Depending on the purpose of the test, the fabrics may subsequently be washed and dried.

The test and control fabrics are cut into 20cm x 20cm squares for testing.

Stage 2 is the carrying out of the test. A team of three Caucasian female assessors of minimum age 20 years is selected to carry out olfactory evaluation on the basis that each is able to rank correctly the odour levels of the series of standard aqueous solutions of isovaleric acid listed below, and each is able to assign a numerical score, corresponding to the odour intensity of one of these solutions, to the body malodour of a shirt insert after has been worn in the axillary region by a male subject for a standard period of time.

A panel of 40 human subjects for use in the test is assembled from Caucasian male subjects of age within the range of from 20 to 55 years. By screening, subjects are chosen who develop axillary body malodour that is not unusually strong and who do not develop a stronger body malodour in one axilla compared with the other. Subjects who develop unusually strong body malodour, for example due

to a diet including curry or garlic, are not selected for the panel.

For two weeks before the start of the test, the panel subjects are assigned an unperfumed, non-deodorant soap bar for exclusive use when washing and are denied the use of
5 any other type of deodorant or antiperspirant. At the end of this period, the 40 subjects are randomly divided into two groups of 20.

The "test" and "control" fabric pieces are then tacked into 40 clean cotton or polyester cotton shirts in the underarm
10 region in such a manner that in 20 shirts, the control fabric pieces are attached inside the left underarm region, and the test fabric pieces are attached in the right underarm region. For the remaining 20 shirts, the placing of control and test pieces of fabric is reversed.

15 The shirts carrying the tacked-in fabric inserts are then worn by the 40 panel members for a period of 5 hours, during which time each panellist performs his normal work function without unnecessary exercise.

After this five hour period, the shirts are removed and the
20 inserts detached and placed in polyethylene pouches prior to assessment by the trained panel of assessors.

The malodour intensity of each fabric insert is evaluated

by all three assessors who, operating without knowledge of which inserts are "test" and which are "control" and, without knowing the scores assigned by their fellow assessors, sniff each fabric piece and assign to it a score corresponding to the strength of the odour on a scale from 0 to 5, with 0 representing no odour and 5 representing very strong odour.

Standard aqueous solutions of isovaleric acid which correspond to each of the scores 1, 2, 3, 4 and 5 are provided for reference to assist the assessors in the malodour evaluation. These are shown below:

SCORE	ODOUR LEVEL	CONCENTRATION OF AQUEOUS ISOVALERIC ACID (ML/L)
0	NO ODOUR	0
1	SLIGHT	0.013
2	DEFINITE	0.053
3	MODERATE	0.22
4	STRONG	0.87
5	VERY STRONG	3.57

The scores recorded by each assessor for each fabric piece are averaged. The average score of the "test" fabric pieces is deducted from the average score of the "untreated" control fabric pieces to give a Malodour Reduction Value.

As a check that the selection of panel subjects is satisfactory for operation of the test, the average score

with unperfumed fabric pieces should be between 2.5 and 3.0.

The higher the Malodour Reduction Value of at least 0.50, or 0.70, or 1.00. The higher the minimum value, the more effective is the perfume as a deodorant as recorded by the assessors in the Malodour Reduction Value Test. It has also been noted that consumers, who are not trained assessors, can detect by self-assessment a noticeable reduction in malodour on soiled fabric such as shirts and underclothes where the Malodour Reduction Value is at least 0.30, so the higher the Malodour Reduction Value above this figure, the more noticeable is the deodorant effect.

Selection of a combination of fragrance materials to give a deodorant effect is explained in patents such as US-A-430679 referred to earlier. Further systems of selection are given in US-A-5482635 and US-A-5554588 also mentioned above.

Such selections can be carried out using materials with preferred values of partition coefficient and Kovats index as discussed above.

US-A-5501805 describes perfume compositions made from a combination of fragrance materials, where the composition is a deodorant perfume yet has a relatively low odour. Such "low-odour" deodorant perfumes may be used in the present

invention.

When this test is used to assess the deodorant effectiveness of a perfume composition, applied to fabric in accordance with this invention, the test fabric is a blend of 95% by weight cotton and 5% by weight spandex
5 fibres. The control fabric is 100% cotton. The test and control fibres are selected to be similar in other respects, in particular to have the same weight per unit area.

The test fabric is subjected to treatment with a fabric
10 finishing liquor, containing perfume, so as to apply 0.5% of the perfume, by weight of the fabric. The control fabric is treated similarly, but without perfume.

The test and control fabrics are not subsequently washed before testing.

15 However, this test procedure can be operated in other ways. To demonstrate the higher deposition of perfume on spandex fibres, the control and test fabrics are both treated with the same fabric finishing liquor containing perfume. To isolate the deodorant effect of the perfume, the test and
20 control fabrics can be the same, but no perfume is present in the liquor used to treat the control fabric.

Example 1

This model experiment demonstrates perfume deposition on spandex fibres. A mixture of perfume ingredients was prepared and added to an unperfumed, but otherwise conventional, laundry detergent powder, to provide a
5 perfume concentration of 0.5% by weight.

The perfumed powder was used to wash test cloths which had not previously been treated with any perfume. These were either all cotton, or 95% cotton with 5% spandex. After washing, the cloths were rinsed and then line dried
10 overnight.

The perfume was extracted from the dry cloths with organic solvent, and the content of the perfume ingredients in the solvent extracts was determined by gas chromatography. If the concentration of an ingredient extracted from the
15 spandex containing cloth was greater than from the all-cotton cloth by a factor of 5 to 20, the result was coded as a medium enhancement(M). If the concentration was greater by 20 or more, it was coded high(H) and if less than 5 or not measurable, it was coded(L). The results
20 obtained were as follows:

Ingredient	K*	logP* *	Enhancement	Category
Boisambrene Forte	1714	5.5	M	B
benzyl acetone	1206	2.0	M	-
citronellol	1209	3.6	H	A'
2,6-dimethyl- heptan-2-ol	975	2.9	L	-
jasmacyclene	1394	2.9	H	B'
methyl salicylate	1167	2.3	L	-
2-phenylethanol	1087	1.4	L	-
terpinyl acetate	1331	4.0	H	B'
tetrahydrogeraniol	1180	3.6	H	A'
tetrahydrolinalol	1083	3.5	H	A'
Tonalid	1840	6.4	M	B
yara	1416	3.2	H	B'

* Measured on OV-1 polydimethylsiloxane (Ohio Valley) as
stationary phase using capillary gc

** Measured or estimated using 'logP' software from ACD
Inc.

Example 2

Cloth composed of 90% cotton, 10% spandex was treated in a fabric finishing step, using conventional equipment. Other cloth, consisting of cotton only, was treated in the same way. In both cases, the fabric finishing treatment was

5 carried out for a period of 20 minutes, using an MCS jet machine (Urgano, Italy). The finishing liquors were applied applied at a liquor to cloth ratio of 20:1, at 40°C with a pH of 5.5. These liquors all contained Ceranine HCS (a finishing agent made by Sandoz) at a concentration such

10 that this agent was applied to fabric at 1% by weight of the fabric.

The treatment liquors also contained perfume at varying concentrations, so that this was applied to the fabric at concentrations of 0.01%, 0.1% and 1% by weight of the

15 fabric. Liquor without perfume was used to provide a control.

The perfume was used, designated "perfume U" contained 33.5% (by weight of the perfume composition) of fragrance materials in category A above, all of which had Kovats

20 indices of 1050 to 1600 and therefore all fell within the narrower category A'. The perfume also contained 41.1% (by weight of the perfume composition) of materials in category B. These were made up from 26.7% with Kovats index over 1600 and 14.4% with Kovats index of 1300 to 1600, so as to

fall within category B'.

The cloths were then washed repeatedly, using a commercial detergent powder which included a different perfume. The washes were carried out using a Miele washing machine on its "Quickwash" programme at 40°C. 110gm of detergent powder was used for each wash. The fabric was rinsed three times after each wash and tumble dried.

The dried cloths were examined by a panel of expert assessors of perfume intensity. This was to determine the intensity of perfume on the fabric, but not its deodorant property. The results obtained were as follows:

	Odour Evaluation Scores					
	100% cotton			90% cotton and 10% spandex		
Washes:	1	3	5	1	3	5
Perfume U						
0%	2.0	2.4	2.8	4.0	4.4	4.6
1.0%	6.4	3.6	<3	16.0	14.0	12.8
0.1%	3.6	<3	<3	10.4	9.7	9.0
0.01%	3.2	<3	<3	8.8	8.0	7.2

It can be seen that the cloths which were not perfumed in the finishing treatment took up perfume in the first wash and this perfume built up slowly in subsequent washes. The quantity of perfume taken up was greater on the cloth which included spandex fibres.

The cloths containing spandex fibres which were perfumed during the finishing treatment had a much higher level of perfume on them after one wash than the 100% cotton cloths. Even after five washes, the intensity of perfume on them exceeded the intensity of perfume on the 100% cotton cloths after one wash, and on the cloths which had not been perfumed prior to the first wash. Thus the spandex fibres were providing enhanced retention of perfume as well as enhanced deposition.

The 100% cotton cloths which had been perfumed during the finishing treatment were assessed again after 3 and 5 washes. The results showed that the level of intensity of the perfume was less than that observed after 1 wash but also showed that the olfactive differences between perfume U used in the finishing treatment and the perfume present in the washing powder was confusing the panellists.

Example 3

Two deodorant perfumes were used in treatment of cloths by a finishing process as in Example 2.

Perfume L contained materials in the above categories as follows:

Category A: 30 wt% (all with Kovats index above 1600)

Category B: 68.5 wt% (13% with Kovats index 1300-1600 and therefore within category B', and 55.5 wt% with Kovats index above 1600).

Perfume M contained materials in the above categories as follows:

5 Category A: 24.9 wt% (16.3 wt% with Kovats index 1050-1600 and therefore within category A', and 8.6 wt% with Kovats index above 1600)

Category B: 55.3 wt% (8.6 wt% with Kovats index 1300-1600 and therefore within category B', and 46.7
10 wt% with Kovats index above 1600).

The test cloths were: 100% cotton, 90% cotton with 10% spandex, 95% cotton with 5% spandex, 100% nylon and 82% nylon with 18% spandex. Perfume was used at a concentration of 0.5% based on the weight of the fabric. The treated
15 cloths were tested for Malodour reduction in the test described earlier. The control cloths were 100% cotton, which had been subjected to the same finishing treatment, but without perfume in that finishing treatment. The results are set out in the following tables, which show
20 substantial enhancements of malodour inhibition when fabrics containing spandex fibres were used.

Test 1:

Fabric		perfume M (% by weight of fabric)	Malodour score	Malodour reduction	Malodour reduction as % of control
other fibre	spandex				
90% cotton	10%	0.5%	1.19	1.46	55%
100% cotton	0	0.5%	1.92	0.73	27.4%
82% nylon	18%	0.5%	1.00	1.65	62.1%
100% nylon	0	0.5%	1.97	0.68	25.5%
100% cotton (control)	0	0	2.65		

Notes: % malodour reduction calculated as $100\% \times (\text{control score} - \text{sample score}) / \text{control score}$

Statistical calculation showed that a difference in malodour reduction of 6.9% was significant at 95% level of confidence.

Test 2:

	Fabric		perfume (% by weight of fabric)	Malodour score	Malodour reduction	Malodour reduction as % of control
	other fibre	spandex				
5	95% cotton	5%	0.5% L	1.15	1.22	51.5%
	95% cotton	5%	0.5% M	1.29	1.08	45.6%
10	100% cotton (control)	0	0	2.37		

Note: Statistical calculation showed that a difference in malodour reduction of 6.3% was significant at 95% level of confidence.

In test 1, malodour scores on 100% cotton fabric, with and without perfume, demonstrate a malodour reduction value of 0.73 attributable to the perfume. A similar malodour reduction value was observed when the test cloth was 100% nylon.

When spandex fibre was incorporated, the malodour reduction increased greatly, showing that increased deposition of perfume on spandex fibres compared with other fibres also provides an increased deodorant efficiency.

In test 2, similar high values of malodour reduction were obtained when either perfume L or perfume M was used, in test fabrics with 5% by weight spandex fibres.

Claims

1. A method of treating textile which is yarn or unworn fabric containing spandex fibres, comprising contacting the textile with a perfume composition which is a mixture of fragrance materials, so that
5 fragrance materials are deposited on the textile.

2. A method according to claim 1 wherein the textile contains from 0.5 wt % to 50 wt % spandex fibres and the amount of fragrance materials deposited on the textile is from 0.001% to 1% by weight of the
10 textile.

3. A method according to claim 1 or claim 2 wherein the perfume composition contains at least 50%, by weight of the perfume composition, of fragrance materials selected from:

15 Category A) hydroxylic materials which are alcohols, phenols or salicylates, with an octanol/water partition coefficient (P) whose common logarithm ($\log_{10}P$) is 2.5 or greater, and a gas chromatographic Kovats index (as determined on polydimethylsiloxane
20 as non-polar stationary phase) of at least 1050, and

Category B) esters, ethers, nitriles, ketones or aldehydes, with an octanol/water partition

coefficient (P) whose common logarithm ($\log_{10}P$) is 2.5 or greater, and a gas chromatographic Kovats index (as determined with polydimethylsiloxane as non-polar stationary phase) of at least 1300.

4. A method according to claim 3 wherein the perfume composition contains at least 10%, by weight of the perfume composition, of fragrance materials selected from:

Category A') hydroxylic materials which are alcohols, phenols or salicylates, with an octanol/water partition coefficient (P) whose common logarithm ($\log_{10}P$) is 2.5 or greater, and a gas chromatographic Kovats index (as determined on polydimethylsiloxane as non-polar stationary phase) lying within the range 1050 to 1600, and

Category B') esters, ethers, nitriles, ketones or aldehydes, with an octanol/water partition coefficient (P) whose common logarithm ($\log_{10}P$) is 2.5 or greater, and a gas chromatographic Kovats index (as determined on polydimethylsiloxane as non-polar stationary phase) lying within the range 1300 to 1600.

5. A method according to claim 3 wherein the perfume composition contains at least 70%, by weight of the

perfume composition, of fragrance materials selected from categories A and B, which materials include at least 25% by weight of the perfume composition, of fragrance materials from the categories A' and B' defined in claim 4.

- 5 6. A method according to claim 5 wherein the perfume composition contains at least 80%, by weight of the perfume composition, of fragrance materials selected from categories A and B, which materials include at least 40%, by weight of the perfume composition, from categories A' and B'.
- 10
7. A textile which is yarn or unworn fabric and which contains spandex fibres, having fragrance materials deposited on the textile.
8. A textile according to claim 7 which contains from 0.5 to 50 wt % spandex fibres and the amount of fragrance materials deposited thereon is from 0.001% to 1% by weight of the textile.
- 15
9. A textile according to claim 7 or claim 8 which is yarn, a fabric web or length thereof, or an unworn garment.
- 20
10. Use of a perfume composition which is a mixture of fragrance materials in the treatment of yarn or

unworn fabrics containing spandex and other fibres,
to deposit fragrance materials on the spandex
fibres.

W:\WORK\MFF\UNILEVER\C97082\C97082F\DRAFTSPC.1